

Effect of Phytocompounds from the Heartwood of *Acacia confusa* on Inflammatory Mediator Production

JYH-HORNG WU,[†] YU-TANG TUNG,[‡] SHIH-CHANG CHIEN,[§] SHENG-YANG WANG,[†]
 YUEH-HSIUNG KUO,^{§,¶,||} LIE-FEN SHYUR,^{§,*} AND SHANG-TZEN CHANG^{‡,*}

Department of Forestry, National Chung-Hsing University, Taichung 402, Taiwan; School of Forestry and Resource Conservation and Department of Chemistry, National Taiwan University, Taipei 106, Taiwan, Agricultural Biotechnology Research Center, Academia Sinica, Taipei 115, Taiwan, and Tsuzuki Institute for Traditional Medicine, College of Pharmacy, China Medical University, Taichung 404, Taiwan

Acacia confusa Merr. (Leguminosae) is traditionally used as a medicinal plant in Taiwan. In the present study, anti-inflammatory activity of extracts from the heartwood of *A. confusa* were investigated for the first time. Results demonstrated that ethanolic extracts of *A. confusa* heartwood strongly suppressed NO production in lipopolysaccharide (LPS)-activated RAW 264.7 macrophages. Among all fractions derived from ethanolic extracts, the EtOAc fraction exhibited the best inhibitory activity. Following column chromatography and reverse-phase high-performance liquid chromatography, 13 specific phytocompounds including 5 new flavonoids (i.e., 7,8,3',4'-tetrahydroxy-4-methoxyflavan-3-ol, 7,8,3',4'-tetrahydroxyflavone, 7,8,3'-trihydroxy-3,4'-dimethoxyflavone, 7,3',4'-trihydroxyflavone, and 7,3',4'-trihydroxy-3-methoxyflavone) were isolated and identified from the EtOAc fraction. In addition, melanoxetin (3,7,8,3',4'-pentahydroxyflavone), a major compound in the EtOAc fraction, markedly suppressed LPS-induced NO and prostaglandin E₂ (PGE₂) production. Moreover, melanoxetin completely suppressed gene expression of inducible NO synthase (iNOS) and cyclooxygenase-2 (COX-2) at 50 and 100 μM, respectively. This is the first report to identify the inhibitory bioactivities of melanoxetin on iNOS and COX-2.

KEYWORDS: *Acacia confusa*; melanoxetin; nitric oxide; prostaglandin E₂; inducible nitric oxide synthase; cyclooxygenase-2

INTRODUCTION

Many plant infusions frequently used in domestic medicine have anti-inflammatory and pharmacological properties ascribed to the presence of phenolic compounds, especially flavonoids. Flavonoids are polyphenolics occurring in a wide range of plants, with over 8000 chemically distinct compounds identified (1), and are plentiful in the heartwood of several *Acacia* species (2, 3). Secondary metabolites of the genus *Acacia* have been characterized, and this has been thoroughly reviewed by Seigler (4). However, little information is available concerning the chemical compositions and biological activities of *Acacia confusa*.

A. confusa, a species indigenous to Taiwan, is widely distributed over the hills and lowlands of Taiwan and is traditionally used as a medicinal plant (5). In a previous study, we found that the ethanolic extracts of the heartwood not only contain a large amount of polyphenols but also show an excellent antioxidant activity (6). However, the application of this plant to inflammatory disease has not been supported by pharmacological investigations to date. Numerous studies have revealed that improper up-regulation of inducible nitric oxide synthase (iNOS) and/or cyclooxygenase-2 (COX-2) was associated with the pathophysiological features of certain types of human cancers as well as inflammatory disorders (7–9). Therefore, the identification of phytochemicals that can suppress or down-regulate the functions of iNOS and/or COX-2 may lead to the discovery of important anti-inflammatory therapeutics.

The aim of this study was to investigate the inhibitory effects of natural phytochemicals, particularly melanoxetin, from *A. confusa* heartwood on NO and prostaglandin E₂ (PGE₂) production in RAW 264.7 macrophages. In addition, we sought underlying mechanisms leading to the suppression of NO and PGE₂ generation after melanoxetin treatment.

* Authors to whom correspondence should be addressed [telephone 886-2-33664626; fax 886-2-23654520; e-mail (S.-T.C.) peter@ntu.edu.tw; (L.-F.S.) lfshyur@ccvax.sinica.edu.tw].

[†] National Chung-Hsing University.

[‡] School of Forestry and Resource Conservation, National Taiwan University.

[§] Academia Sinica.

[¶] Department of Chemistry, National Taiwan University.

^{||} China Medical University.




Figure 1. Chemical structures of phytochemicals isolated from the EtOAc soluble fraction of *A. confusa* heartwood: **1**, 3,4-dihydroxybenzoic acid; **2**, 7,8,3',4'-tetrahydroxy-4-methoxyflavan-3-ol; **3**, 7,8,3',4'-tetrahydroxyflavanone; **4**, 3,4-dihydroxybenzoic acid methyl ester; **5**, melanoxetin; **6**, 7,8,3',4'-tetrahydroxyflavone; **7**, transilitin; **8**, 3,4-dihydroxybenzoic acid ethyl ester; **9**, 3,4,2',3',4'-pentahydroxy-*trans*-chalcone; **10**, 3,7,8,3'-tetrahydroxy-4'-methoxyflavone; **11**, 7,8,3'-trihydroxy-3,4'-dimethoxyflavone; **12**, 7,3',4'-trihydroxyflavone; **13**, 7,3',4'-trihydroxy-3-methoxyflavone.

MATERIALS AND METHODS

Reagents. Lipopolysaccharide (LPS), aspirin, sodium pyruvate, streptomycin, *N*-(1-naphthyl)ethylenediamine dihydrochloride, sulfanilamide, dimethyl sulfoxide (DMSO), and quercetin were purchased from Sigma Chemical Co. (St. Louis, MO). The other chemicals and solvents were of the highest quality available.

Extraction and Purification of Phenolic Compounds from the Heartwood of *A. confusa*. The heartwood of *A. confusa* Merr. (40-year-old) was sampled from the experimental forest of National Taiwan University in Nan-Tou County, Taiwan. The species was identified by Sheng-You Lu of the Taiwan Forestry Research Institute, and a voucher specimen (AC001) was deposited at the School of Forestry and Resource Conservation, National Taiwan University. The dried samples (1.65 kg) were cut into small pieces and soaked in 70% ethanol at ambient temperature for 7 days. The extract was decanted, filtered under vacuum, concentrated in a rotary evaporator, and then lyophilized. The resulting crude extract (135 g) was fractionated successively with *n*-hexane, ethyl acetate (EtOAc), butanol (BuOH), and water to yield soluble fractions of *n*-hexane (1 g), EtOAc (84 g), BuOH (20 g), and H₂O (13 g). Thirteen phytochemicals, as shown in **Figure 1**, were isolated and purified from the EtOAc fraction by semipreparative HPLC on a model PU-980 instrument (Jasco) with a 250 × 10.0 mm inside diameter, 5 μm Luna RP-18 column (Phenomenex, Torrance, CA). The mobile phase was solvent A (100% MeOH) and solvent B (ultrapure water). Elution conditions were 0–20 min of 40–60% A to B (linear gradient) and 20–28 min of 60–100% A to B (linear gradient) at a flow rate of 4 mL/min using a Jasco MD-910 photo diode array detector at 280 or 370 nm wavelength. Electron-impact mass spectrometry (EIMS) and high-resolution electron-impact mass spectrometry (HREIMS) data were collected using Finnigan MAT-95S mass spectrometry, and NMR spectra were recorded by a Bruker Avance 500 MHz FT-NMR spectrometer.

7,8,3',4'-Tetrahydroxy-4-methoxyflavan-3-ol (2): light yellow amorphous solid; mp 106–108 °C; UV (MeOH), λ_{max}, nm (log ε), 283.5 (3.82); IR (KBr), ν_{max}, cm⁻¹, 3338, 1613, 1517; HREIMS, *m/z* 320.0898, [M]⁺ calcd 320.0896 for C₁₆H₁₆O₇; ¹H NMR (500 MHz, CD₃OD), δ 7.06 (d, 1, *J* = 1.7 Hz, H-2'), 6.89 (dd, 1, *J* = 8.1, 1.7 Hz,

H-6'), 6.79 (d, 1, *J* = 8.1 Hz, H-5'), 6.69 (d, 1, *J* = 8.3 Hz, H-5), 6.46 (d, 1, *J* = 8.3 Hz, H-6), 4.99 (br s, 1, H-2), 4.09 (d, 1, *J* = 2.8 Hz, H-4), 3.96 (d, 1, *J* = 2.8 Hz, H-3), 3.47 (s, 3, 4-OCH₃); ¹³C NMR (125 MHz, CD₃OD), δ 147.3 (C-7), 146.1 (C-3', C-4'), 145.4 (C-9), 133.9 (C-8), 131.4 (C-1'), 123.3 (C-5), 119.8 (C-6'), 116.0 (C-5'), 115.8 (C-2'), 112.6 (C-10), 109.2 (C-6), 78.7 (C-4), 76.8 (C-2), 69.5 (C-3), 56.5 (4-OCH₃).

7,8,3',4'-Tetrahydroxyflavone (6): light yellow amorphous solid; mp 226–228 °C; UV (MeOH), λ_{max}, nm (log ε), 265 (4.16), 345 (4.13); IR (KBr), ν_{max}, cm⁻¹, 3226, 1640, 1613, 1574, 1517; HREIMS, *m/z* 286.0473, [M]⁺ calcd 286.0477 for C₁₅H₁₀O₆; ¹H NMR (500 MHz, CD₃OD), δ 7.51 (d, 1, *J* = 8.7 Hz, H-5), 7.51 (d, 1, *J* = 8.9 Hz, H-6'), 7.51 (s, 1, H-2'), 6.94 (d, 1, *J* = 8.7 Hz, H-6), 6.91 (d, 1, *J* = 8.9 Hz, H-5'), 6.62 (s, 1, H-3); ¹³C NMR (125 MHz, CD₃OD), δ 180.8 (C-4), 166.0 (C-2), 152.3 (C-7), 150.8 (C-4'), 148.5 (C-9), 147.0 (C-3'), 134.6 (C-8), 124.3 (C-1'), 118.0 (C-10), 116.8 (C-5), 116.7 (C-5'), 115.0 (C-6, C-6'), 114.4 (C-2'), 104.7 (C-3).

7,8,3'-Trihydroxy-3,4'-dimethoxyflavone (11): light yellow amorphous solid; mp 211–213 °C; UV (MeOH), λ_{max}, nm (log ε), 259.5 (4.33), 346.5 (4.25); IR (KBr), ν_{max}, cm⁻¹, 3184, 1660, 1602, 1567, 1513; HREIMS, *m/z* 310.0734, [M]⁺ calcd 310.0739 for C₁₇H₁₄O₇; ¹H NMR (500 MHz, CD₃OD), δ 7.77 (dd, 1, *J* = 8.6, 2.1 Hz, H-6'), 7.74 (d, 1, *J* = 2.1 Hz, H-2'), 7.54 (d, 1, *J* = 8.8 Hz, H-5), 7.08 (d, 1, *J* = 8.6 Hz, H-5'), 6.94 (d, 1, *J* = 8.8 Hz, H-6), 3.94 (s, 3, 4'-OCH₃), 3.77 (s, 3, 3-OCH₃); ¹³C NMR (125 MHz, CD₃OD), δ 177.0 (C-4), 157.5 (C-2), 152.0 (C-7), 151.6 (C-4'), 147.7 (C-9), 147.6 (C-3'), 141.2 (C-3), 134.4 (C-8), 124.8 (C-1'), 122.4 (C-6'), 118.6 (C-10), 116.8 (C-5), 116.5 (C-2'), 115.2 (C-6), 112.3 (C-5'), 60.4 (3-OCH₃), 56.4 (4'-OCH₃).

7,3',4'-Trihydroxyflavone (12): light yellow amorphous solid; mp 294–296 °C; UV (MeOH), λ_{max}, nm (log ε), 235.5 sh (4.27), 316 sh (4.13), 344.5 (4.25); IR (KBr), ν_{max}, cm⁻¹, 3207, 1660, 1621, 1567, 1513; HREIMS, *m/z* 270.0498, [M]⁺ calcd 270.0528 for C₁₅H₁₀O₅; ¹H NMR (500 MHz, CD₃OD), δ 7.96 (d, 1, *J* = 8.7 Hz, H-5), 7.40 (d, 1, *J* = 8.9 Hz, H-6'), 7.39 (s, 1, H-2'), 6.95 (s, 1, H-8), 6.92 (d, 1, *J* = 8.7 Hz, H-6), 6.90 (d, 1, *J* = 8.9 Hz, H-5'), 6.62 (s, 1, H-3); ¹³C NMR (125 MHz, CD₃OD), δ 180.3 (C-4), 166.1 (C-2), 165.0 (C-7), 159.7 (C-9), 150.8 (C-4'), 147.1 (C-3'), 127.7 (C-5), 124.0 (C-1'), 120.2 (C-6'), 117.2 (C-10), 116.8 (C-5'), 116.3 (C-6), 114.1 (C-2'), 105.2 (C-3), 103.5 (C-8).

7,3',4'-Trihydroxy-3-methoxyflavone (13): light yellow amorphous solid; mp 225–227 °C; UV (MeOH), λ_{max}, nm (log ε), 242 sh (4.21), 321 sh (4.07), 349.5 (4.17); IR (KBr), ν_{max}, cm⁻¹, 3192, 1660, 1605, 1563, 1505; HREIMS, *m/z* 300.0625, [M]⁺ calcd 300.0634 for C₁₆H₁₂O₆; ¹H NMR (500 MHz, CD₃OD), δ 7.99 (d, 1, *J* = 8.7 Hz, H-5), 7.64 (d, 1, *J* = 2.2 Hz, H-2'), 7.55 (dd, 1, *J* = 8.4, 2.2 Hz, H-6'), 6.92 (d, 1, *J* = 8.7 Hz, H-6), 6.91 (d, 1, *J* = 8.4 Hz, H-5'), 6.91 (s, 1, H-8), 3.76 (s, 3, 3-OCH₃); ¹³C NMR (125 MHz, CD₃OD), δ 176.5 (C-4), 164.8 (C-7), 158.7 (C-9), 157.9 (C-2), 149.8 (C-4'), 146.5 (C-3'), 141.2 (C-3), 127.8 (C-5), 123.3 (C-1'), 122.3 (C-6'), 117.7 (C-10), 116.5 (C-2'), 116.4 (C-6), 116.2 (C-5'), 103.1 (C-8), 60.3 (3-OCH₃).

Cell Culture. The murine monocyte/macrophage cell line RAW 264.7 was obtained from the Culture Collection and Research Center (CCRC), HsinChu, Taiwan. RAW 264.7 cells were cultured in 75 or 150 cm² plastic flasks with Dulbecco's modified Eagle's medium (DMEM, Gibco/BRL, Grand Island, NY) supplemented with 1 mM sodium pyruvate, penicillin (100 units/mL), streptomycin (100 μg/mL), and 10% heat-inactivated fetal bovine serum (Gibco/BRL). Cells were incubated at 37 °C in a humidified atmosphere of 5% CO₂.

Measurement of NO and PGE₂. RAW 264.7 cells were seeded in 96-well plates at 2 × 10⁵ cells/well and grown for 2 h for adherence. The adhered cells were incubated for 24 h, with or without 1 μg/mL of LPS, in the absence or presence of the test extracts or compounds (DMSO as a vehicle). The nitrite concentration in the culture medium was measured as an indicator of NO production according to the Griess reaction (10). Briefly, 100 μL of culture supernatant or sodium nitrite standard underwent reaction with an equal volume of Griess reagent [a mixture of 0.1% (w/v) *N*-(1-naphthyl)ethylenediamine dihydrochloride in water and 1% (w/v) sulfanilamide in 5% (v/v) phosphoric acid] in a 96-well plate. After 10 min at ambient temperature, the absorbance




Figure 2. Effect of ethanolic extract and its derived soluble fractions of *A. confusa* on the nitric oxide production in lipopolysaccharide (LPS)-stimulated RAW 264.7 macrophage cells: **(A)** effect of ethanolic extract on LPS-induced nitric oxide production; **(B)** cytotoxicity of ethanolic extract on RAW 264.7 cells in the presence of LPS, measured by the MTT assay; **(C)** effect of various fractions of ethanolic extract on LPS-induced nitric oxide production; **(D)** cytotoxicity of various fractions of ethanolic extract on cells in the presence of LPS, measured by the MTT assay. Results are mean \pm SD ($n = 3$). Different letters indicate significant differences among groups ($p < 0.05$).




Figure 3. HPLC chromatogram of EtOAc fraction from the heartwood of *A. confusa*.

at 550 nm was measured by use of a microtitration plate reader (Labsystems, Helsinki, Finland).

For PGE₂ determination, RAW 264.7 cells were seeded in 96-well plates at 1×10^4 cells/well and incubated for 18 h. Cells were pretreated with 500 μ M of aspirin for 3 h to inactivate endogenous cyclooxygenase-1 (COX-1) based on the method reported by Hwang et al. (11). Then, cells were washed twice with phosphate-buffered saline (PBS) and further incubated for 16 h in fresh DMEM with or without 1 μ g/mL of LPS in the absence or presence of the test samples. After incubation, supernatants were collected to measure PGE₂ concentration with monoclonal antibody by enzyme-linked immunosorbent assay (ELISA) as specified by the manufacturer (Cayman Chemical, Ann Arbor, MI).

Cell Viability (MTT Assay). Cell respiration as an indicator of cell viability was determined on the basis of mitochondrial-dependent

reduction of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) to formazan (12). After culture supernatants were collected for NO or PGE₂ measurement, culture wells were washed once with 1 mL of PBS to remove the remaining test samples. Cells were then incubated with 0.5 mg/mL MTT for 4 h at 37 °C in a 5% CO₂ incubator. The medium was aspirated, and the insoluble formazan product was dissolved in DMSO (250 μ L) for 2 h in the dark. The extent of MTT reduction was quantified by measuring the absorbance at $\lambda = 570$ nm.

RT-PCR Analysis. RAW 264.7 cells were seeded in six-well plates at 3×10^6 cells/well. The cells were treated with various concentrations of test compounds for 1 h and then incubated for 6 h in fresh DMEM with or without 1 μ g/mL of LPS. After three washings with ice-cold PBS, the cells were harvested, and total RNA was extracted with use of TRIzol reagent (Life Technologies, Gaithersburg, MD) as specified by the manufacturer. For each RT-PCR reaction, 2 μ g of total RNA was used to synthesize first-strand cDNA with Superscript reverse transcriptase (Life Technologies). Reverse transcriptase-generated cDNA encoding for murine iNOS and COX-2 were amplified by PCR. Oligonucleotide primers for iNOS and COX-2 were 5'-CAGAAGCA-GAATGTGACCATC-3' (sense), 5'-CTTCTGGTTCGATGTCATGAGC-3' (antisense); and 5'-ACTCACTCAGTTTGTGAGTCATTC-3' (sense), 5'-TTTGATTAGTACTGTAGGGTTAATG-3' (antisense), respectively. The cDNA sequence of glyceraldehyde-3-phosphate dehydrogenase (GA3PDH) as an internal control was also amplified using the following primers: 5'-CCATCAATGACCCCTTCATTGACC-3' (sense) and 5'-GAAGGCCATGCCAGTGAGCTCC-3' (antisense). For PCR amplification the following conditions were used: for iNOS and GA3PDH, 94 °C for 2 min for one cycle and then 94 °C for 1 min, 55 °C for 30 s, and 72 °C for 1 min for 30 cycles; for COX-2, 94 °C for 2 min for one cycle and then 94 °C for 15 s, 55 °C for 1 min, and 72 °C for 1 min for 30 cycles. The amplified PCR products were analyzed with 2% agarose gel electrophoresis and visualized with ethidium bromide staining.

Table 1. ^{13}C NMR Spectral Data and HMBC Correlations of Five New Phytocompounds Isolated from the Heartwood of *A. confusa* (CD_3OD)

no.	2		6		11		12		13	
	δ_{C}	HMBC	δ_{C}	HMBC	δ_{C}	HMBC	δ_{C}	HMBC	δ_{C}	HMBC
2	76.8	H-4 H-2' H-6'	166.0	H-3	157.5	H-2' H-6'	166.1	H-3 H-2' H-6'	157.9	H-2' H-6'
3	69.5		104.7		141.2	OCH ₃	105.2		141.2	OCH ₃
4	78.7	H-5 OCH ₃	180.8	H-5	177.0	H-5	180.3	H-3 H-5	176.5	H-5
5	123.3	H-4	116.8		116.8		127.7		127.8	
6	109.2		115.0		115.2		116.3	H-8	116.4	H-8
7	147.3	H-5	152.3	H-5	152.0	H-5 H-6	165.0	H-5 H-8	164.8	H-5 H-8
8	133.9	H-6	134.6	H-6	134.4	H-6	103.5	H-6	103.1	H-6
9	145.4	H-4 H-5	148.5	H-5	147.7	H-5	159.7	H-5 H-8	158.7	H-5 H-8
10	112.6	H-3 H-6	118.0	H-3 H-6	118.6	H-6	117.2	H-3 H-6	117.7	H-6 H-8
1'	131.4	H-2 H-5'	124.3	H-3 H-5'	124.8	H-5'	124.0	H-5'	123.3	H-2' H-5'
2'	115.8	H-2 H-6'	114.4	H-6'	116.5	H-6'	114.1	H-6'	116.5	H-6'
3'	146.1	H-5'	147.0	H-2' H-5'	147.6	H-5'	147.1	H-2' H-5'	146.5	H-2' H-5'
4'	146.1	H-2' H-6'	150.8	H-2' H-6'	151.6	H-2' H-6' OCH ₃	150.8	H-2' H-5' H-6'	149.8	H-2' H-5' H-6'
5'	116.0		116.7		112.3		116.8		116.2	
6'	119.8	H-2 H-2'	115.0	H-2'	122.4	H-2'	120.2	H-2'	122.3	H-2'

Statistical Analyses. All results are expressed as mean \pm SD ($n = 3$). Statistical analyses were performed using ANOVA with Scheffé's test, and $p < 0.05$ was considered to be significant.

RESULTS AND DISCUSSION

Effects of Phytochemicals from *A. confusa* Heartwood on NO Production in RAW 264.7 Cells. NO is a critical signaling molecule produced at inflammatory sites by iNOS, which is often expressed in response to LPS, interferon- γ , and various proinflammatory cytokines (13). Several physiological events by which intracellular elevated NO can exert genotoxic effects after reacting with oxygen include formation of carcinogenic *N*-nitroso compounds, DNA strand breakage, and nitrosative deamination of DNA bases (14, 15). Thus, maintenance of a tight regulation of NO production is important for human health. In our previous study, the ethanolic extract (6) and phenolic compounds (16) from the heartwood of *A. confusa* exhibited excellent inhibitory activity against reactive oxygen species (ROS). To determine further the effects of those extracts on NO production, different concentrations of test samples were incubated with LPS-activated RAW 264.7 macrophages. As shown in **Figure 2A**, the nitrite level in cultured supernatant of RAW 264.7 cells was markedly elevated, from 0.8 to 16.6 μM , after 24 h of treatment with LPS. An ethanolic extract of *A. confusa* heartwood significantly inhibited LPS-induced NO production in a dose-dependent manner. Relative to LPS treatment only, 28% of NO production was inhibited in the ethanolic extract-treated cells at the concentration of 50 $\mu\text{g}/\text{mL}$. To evaluate whether the inhibition of NO production was possibly caused by the cytotoxic effect of test extracts, the viability of test cells was determined by the MTT assay. **Figure 2B** shows that the ethanolic extract of heartwood shows no significant cytotoxicity to RAW 264.7 cells at all test concentrations. Thus, the inhibition of NO production in LPS-stimulated RAW 264.7 cells by the ethanolic extract of *A. confusa* heartwood was not due to cytotoxicity.

As for the inhibitory effects of various soluble fractions derived from the ethanolic extract on NO production, data in **Figure 2C** show that except for the water soluble fraction, all of the test samples exhibited significant and differential inhibitory activities against NO generation. The IC_{50} values of *n*-hexane, EtOAc and BuOH soluble fractions were 208, 31, and 42 $\mu\text{g}/\text{mL}$, respectively. Of these, the EtOAc fraction exhibited the strongest activity. The addition of 10, 50, 100, and 250 $\mu\text{g}/\text{mL}$ EtOAc extract to LPS-stimulated cells resulted in decreased NO production, by 71, 44, 30, and 22%, respectively, as compared to that of the cells treated with LPS alone. In addition, the MTT assay revealed that concentrations up to 100 $\mu\text{g}/\text{mL}$ produced no cytotoxic effects on cells treated with EtOAc extract (**Figure 2D**).

In our previous study, the EtOAc extract showed a stronger antioxidant activity than other fractions (6). These results indicate that antioxidant and anti-inflammatory activities of ethanolic extract of *A. confusa* heartwood have been effectively enriched in the EtOAc fraction. Moreover, the EtOAc fraction had 84- and 4.2-fold more mass than the *n*-hexane and BuOH fractions, respectively. Thus, it was further investigated in this study for its phytochemical characteristics. **Figure 3** shows the HPLC chromatogram of the EtOAc fraction, from which 13 phenolic compounds (**1–13**) were isolated and identified as 3,4-dihydroxybenzoic acid (**1**), 7,8,3',4'-tetrahydroxy-4-methoxyflavan-3-ol (**2**), 7,8,3',4'-tetrahydroxyflavanone (**3**), 3,4-dihydroxybenzoic acid methyl ester (**4**), melanoxetin (**5**), 7,8,3',4'-tetrahydroxyflavone (**6**), transilitin (**7**), 3,4-dihydroxybenzoic acid ethyl ester (**8**), 3,4,2',3',4'-pentahydroxy-*trans*-chalcone (**9**), 3,7,8,3'-tetrahydroxy-4'-methoxyflavone (**10**), 7,8,3'-trihydroxy-3,4'-dimethoxyflavone (**11**), 7,3',4'-trihydroxyflavone (**12**), and 7,3',4'-trihydroxy-3-methoxyflavone (**13**), respectively. Among these, the ^1H , ^{13}C NMR, and MS spectral data of compounds **1**, **3–5**, and **7–10** were in good agreement with the published values (16–22). Additionally, five flavonoids, including compounds **2**, **6**, **11**, **12**, and **13**, were for the first time isolated

from nature and reported here as new natural products. All ^{13}C signals and heteronuclear multiple bond correlations (HMBC) of these new phytochemicals are shown in **Table 1**. Furthermore, the relative stereochemistry of compound **2** was assigned on the basis of coupling constants and NOESY data. Particularly important was the NOESY correlation seen from H2 (a broad singlet at δ_{H} 4.99) to H3 (δ_{H} 3.96), which placed these two protons in vicinal axial-equatorial orientations. In addition, H2 proton also showed NOESY correlation to H4 (δ_{H} 4.09), and so H4 could be assigned to an axial orientation. Accordingly, this assigned stereochemistry confirmed compound **2** as (2,3-*cis*-3,4-*cis*)-7,8,3',4'-tetrahydroxy-4-methoxyflavan-3-ol.

On the other hand, melanoxetin and transilitin were found as two major flavonoids in the EtOAc fraction. The contents of melanoxetin and transilitin were determined as 33.4 ± 1.1 and 24.5 ± 2.0 $\mu\text{g}/\text{mg}$ of dry weight of EtOAc extract, respectively. The effect of the two specific flavonoids of *A. confusa* on the inhibition of NO generation in LPS-stimulated RAW 264.7 cells was further investigated. Quercetin, one of flavonoids known for its anti-inflammatory activity (23, 24), was used in parallel as a positive control. **Figure 4A** shows that melanoxetin and quercetin strongly inhibited LPS-induced NO production in a concentration-dependent manner, with IC_{50} values of 6.9 and 6.4 μM , respectively. In contrast, transilitin showed poor inhibitory effects ($\text{IC}_{50} > 100$ μM). Except for quercetin, which exhibited a slight cytotoxicity at 100 μM , no effects were observed for melanoxetin and transilitin at the same tested concentrations (**Figure 4B**). This result implies that the 3-methoxyl moiety in the C-ring of transilitin significantly attenuated its inhibitory activity on NO production.

Effect of Melanoxetin on PGE₂ Release from LPS-Activated Macrophages. The activation of macrophages plays a critical role in the inflammatory process by releasing a variety of inflammatory mediators (11). Prostaglandins (PGs) are one of the potent modulators in both inflammation and immune responses. High levels of PGE₂ can affect the activities of several cell types, including neurons, glia, and endothelial cells, and can regulate microglia/macrophage and lymphocyte functions during inflammatory and immune processes (25). In addition, as compared with the surrounding normal tissues, elevated levels of PGs, especially PGE₂, have also been found in many human cancers (26). Thus, phytochemicals that can effectively inhibit the abnormal production of PGE₂ may be potential anti-inflammatory and cancer chemopreventive agents. In the present study, the accumulation of PGE₂ in RAW 264.7 cells was increased from 31.6 to 1383.7 pg/mL by LPS treatment. However, melanoxetin and quercetin inhibited the LPS-induced accumulation of PGE₂ in a dose-dependent manner. **Figure 5** shows that, at a dose of 100 μM , melanoxetin markedly suppressed ($\sim 60\%$) the PGE₂ production in LPS-stimulated RAW 264.7 cells. Quercetin exhibited 98% inhibition of PGE₂ production, but slight cytotoxicity was observed at the same concentration (data not shown). Thus, the hydroxyl moieties in the A-ring of flavonoids play a role and exhibit differential effects on PGE₂ production in macrophages (i.e., the 5,7-dihydroxyl groups in quercetin possess higher activity than the 7,8-dihydroxyl groups in melanoxetin).

Effect of Melanoxetin on iNOS mRNA and COX-2 mRNA Levels in LPS-Activated RAW 264.7 Macrophages. Nitric oxide synthase (NOS) catalyzes NO synthesis in biological systems. Three distinct isoforms of NOS have been isolated, including a neuronal (nNOS), an endothelial (eNOS), and an endotoxin- or cytokine-inducible (iNOS) form (27). Both nNOS and eNOS, often grouped together as constitutive NOS (cNOS), are usually constitutively expressed, and their activities are




Figure 4. Effect of melanoxetin, transilitin, and quercetin on the nitric oxide production of lipopolysaccharide-stimulated RAW 264.7 macrophage cells: (A) concentration-dependent inhibition of nitric oxide production; (B) cytotoxicity of phytochemicals on cells in the presence of LPS, measured by the MTT assay. Results are mean \pm SD ($n = 3$). Different letters indicate significant differences among groups ($p < 0.05$).




Figure 5. Effect of melanoxetin and quercetin on the prostaglandin E₂ production of lipopolysaccharide-stimulated RAW 264.7 macrophage cells. Prostaglandin E₂ in the culture medium was measured as described under Materials and Methods. Results are mean \pm SD ($n = 3$). Different letters indicate significant differences among groups ($p < 0.05$).

regulated by intracellular calcium concentrations via calmodulin. iNOS is usually not detectable in healthy tissues but is expressed after immunological challenge or injury. Enhanced expression of iNOS and its enzymatic activity have been observed in many human tumor tissues and in inflammatory disorders (7–9, 13). In this study, the effect of melanoxetin on iNOS gene expression was also investigated. As shown in **Figure 6A**, unstimulated RAW 264.7 cells (vehicle control) showed barely detectable




Figure 6. Effect of melanoxtetin on the expression of iNOS mRNA (A) and COX-2 mRNA (B) in lipopolysaccharide (LPS)-stimulated RAW 264.7 macrophage cells. Cells were incubated with or without LPS (1 $\mu\text{g/mL}$) in the presence of the indicated concentrations of melanoxtetin.

iNOS mRNA. In contrast, 6 h of incubation with LPS resulted in a large increase in iNOS mRNA expression. However, at 50 μM , melanoxtetin completely suppressed the iNOS mRNA expression. The proposed mechanisms associated with the reduction of NO production are scavenging of NO, suppression of iNOS enzyme activity, inhibition of iNOS gene expression, and/or down-regulation of iNOS enzyme by modulation of enzyme activities related to signal transduction, etc. (28–32). In the present study, melanoxtetin inhibited NO production in macrophages via one or more of these mechanisms.

Cyclooxygenase (COX) is the rate-limiting enzyme in PG synthesis and exists as two isoforms: constitutive (COX-1) and inducible (COX-2) (29). Like iNOS, COX-2 is an important enzyme that mediates inflammatory processes. Multiple lines of compelling evidence support COX-2 playing a role in the

development of tumors (13). Thus, aberrant or excessive expression of iNOS and COX-2 is implicated in inflammatory disorders and the pathogenesis of cancer. Melanoxtetin not only showed strong inhibitory activity on iNOS mRNA expression but also significantly inhibited the COX-2 mRNA expression in LPS-stimulated macrophages. **Figure 6B** shows that melanoxtetin suppressed the LPS-induced COX-2 mRNA expression in a dose-dependent manner. Approximately 5 and 76% reductions were observed at 10 and 50 μM , respectively, as determined by densitometry analysis. At concentrations up to 100 μM , melanoxtetin could completely inhibit the expression of COX-2 mRNA in LPS-stimulated cells. Thus, one of the mechanisms of melanoxtetin inhibition of NO and PGE₂ production in LPS-stimulated macrophages is mediated by the down-regulation of iNOS and COX-2 mRNA expression. Further studies on intracellular signaling cascades leading to COX-2 and iNOS reduction by melanoxtetin are of interest. In addition, *in vivo* pharmacological research on the anti-inflammatory activity of melanoxtetin should also be addressed.

Phytochemicals, especially flavonoids and phenolic acids, are of current interest because of their important biological and pharmacological properties, including reactive oxygen species (ROS) scavenging and anti-inflammatory, anticancer, antimutagenic, and anticarcinogenic activities (33–36). Rice-Evans and Miller (37) have demonstrated the potential role of phytochemicals, including phenylpropanoids, flavonoids, and phenolic acids, as important contributing factors to dietary antioxidant activity. The heartwood of *A. confusa* contains abundant phenolic compounds, especially flavonoids, and shows significant antioxidant and anti-inflammatory activities. Thus, heartwood extracts or the derived phytochemicals from *A. confusa* have great potential to prevent diseases caused by the overproduction of reactive oxygen species and inflammatory disorders.

ACKNOWLEDGMENT

We thank the experimental forest of National Taiwan University for providing *A. confusa* materials, as well as Shou-Ling Huang (Department of Chemistry, National Taiwan University) for NMR spectral analyses.

LITERATURE CITED

- Pietta, P. G. Flavonoids as antioxidants. *J. Nat. Prod.* **2000**, *63*, 1035–1042.
- Clark-Lewis, J. W.; Porter, L. J. Phytochemical survey of the heartwood flavonoids of *Acacia* species from arid zones of Australia. *Aust. J. Chem.* **1972**, *25*, 1943–1955.
- Pietarinen, S. P.; Willför, S. M.; Sjöholm, R. E.; Holmbom, B. R. Bioactive phenolic substances in important tree species. Part 3: Knots and stemwood of *Acacia crassicarpa* and *A. mangium*. *Holzforchung* **2005**, *59*, 94–101.
- Seigler, D. S. Phytochemistry of *Acacia*—sensu lato. *Biochem. Syst. Ecol.* **2003**, *31*, 845–873.
- Kan, W.-S. Leguminosae. In *Manual of Medicinal Plants in Taiwan*; Kan, W.-S., Ed.; National Research Institute of Chinese Medicine: Taipei, Taiwan, 1978; Vol. 2, pp 239–240.
- Chang, S.-T.; Wu, J.-H.; Wang, S.-Y.; Kang, P.-L.; Yang, N.-S.; Shyur, L.-F. Antioxidant activity of extracts from *Acacia confusa* bark and heartwood. *J. Agric. Food Chem.* **2001**, *49*, 3420–3424.
- Thomsen, L. L.; Lawton, F. G.; Knowles, R. G.; Breesley, J. E.; Riveros-Moreno, V.; Moncada, S. Nitric oxide synthase activity in human gynecological cancer. *Cancer Res.* **1994**, *54*, 1352–1354.
- Thomsen, L. L.; Miles, D. W.; Happerfield, L.; Bobrow, L. G.; Knowles, R. G.; Moncada, S. Nitric oxide synthase activity in human breast cancer. *Br. J. Cancer* **1995**, *72*, 41–44.

- (9) Vane, J. R.; Mitchell, J. A.; Appleton, I.; Tomlinson, A.; Bishop, B. D.; Croxtall, J.; Willoughby, D. A. Inducible isoforms of cyclooxygenase and nitric oxide synthase in inflammation. *Proc. Natl. Acad. Sci. U.S.A.* **1994**, *91*, 2046–2050.
- (10) Kim, H.; Lee, H. S.; Chang, K. T.; Ko, T. H.; Baek, K. J.; Kwon, N. S. Chloromethyl ketones block induction of nitric oxide synthase in murine macrophages by preventing activation of nuclear factor- κ B. *J. Immunol.* **1995**, *154*, 4741–4748.
- (11) Hwang, B. Y.; Lee, J.-H.; Koo, T. H.; Kim, H. S.; Hong, Y. S.; Ro, J. S.; Lee, K. S.; Lee, J. J. Furanoligularenone, an eremophilane from *Ligularia fischeri*, inhibits the LPS-induced production of nitric oxide and prostaglandin E₂ in macrophage RAW 264.7 cells. *Planta Med.* **2002**, *68*, 101–105.
- (12) Mosmann, T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J. Immunol. Methods* **1983**, *65*, 55–63.
- (13) MacMicking, J.; Xie, Q. W.; Nathan, C. Nitric oxide and macrophage function. *Annu. Rev. Immunol.* **1997**, *15*, 323–350.
- (14) Iyengar, R.; Stuehr, D. J.; Marletta, M. A. Macrophage synthesis of nitrite, nitrate, and *N*-nitrosamines: precursors and role of the respiratory burst. *Proc. Natl. Acad. Sci. U.S.A.* **1987**, *84*, 6369–6373.
- (15) Nguyen, T.; Brunson, D.; Crespi, C. L.; Penman, B. W.; Wishnok, J. S.; Tannenbaum, S. R. DNA damage and mutation in human cells exposed to nitric oxide in vitro. *Proc. Natl. Acad. Sci. U.S.A.* **1992**, *89*, 3030–3034.
- (16) Wu, J.-H.; Tung, Y.-T.; Wang, S.-Y.; Shyur, L.-F.; Kuo, Y.-H.; Chang, S.-T. Phenolic antioxidants from the heartwood of *Acacia confusa*. *J. Agric. Food Chem.* **2005**, *53*, 5917–5921.
- (17) Flamini, G.; Antognoli, E.; Morelli, I. Two flavonoids and other compounds from the aerial parts of *Centaurea bracteata* from Italy. *Phytochemistry* **2001**, *57*, 559–564.
- (18) Foo, L. Y. Configuration and conformation of dihydroflavonols from *Acacia melanoxylon*. *Phytochemistry* **1987**, *26*, 813–817.
- (19) Miyazawa, M.; Oshima, T.; Koshio, K.; Itsuzaki, Y.; Anzai, J. Tyrosinase inhibitor from black rice bran. *J. Agric. Food Chem.* **2003**, *51*, 6953–6956.
- (20) Malan, E. 7,8,4'-Trihydroxy-3,3'-dimethoxyflavone from the heartwood of *Acacia nigrescens*. *Phytochemistry* **1993**, *33*, 733–734.
- (21) Baderschneider, B.; Winterhalter, P. Isolation and characterization of novel benzoates, cinnamates, flavonoids, and lignans from riesling wine and screening for antioxidant activity. *J. Agric. Food Chem.* **2001**, *49*, 2788–2798.
- (22) Hoffmann, B.; Hölzl, J. Chalcone glucosides from *Bidens pilosa*. *Phytochemistry* **1989**, *28*, 247–249.
- (23) Kim, H. K.; Cheon, B. S.; Kim, Y. H.; Kim, S. Y.; Kim, H. P. Effects of naturally flavonoids on nitric oxide production in the macrophage cell line RAW 264.7 and their structure–activity relationships. *Biochem. Pharmacol.* **1999**, *58*, 759–765.
- (24) Wadsworth, T. L.; Koop, D. R. Effects of the wine polyphenolics quercetin and resveratrol on pro-inflammatory cytokine expression in RAW 264.7 macrophages. *Biochem. Pharmacol.* **1999**, *57*, 941–949.
- (25) Weissmann, G. Prostaglandins as modulators rather than mediators of inflammation. *J. Lipid Mediat.* **1993**, *6*, 275–286.
- (26) Surh, Y.-J.; Chun, K.-S.; Cha, H.-H.; Han, S. S.; Keum, Y.-S.; Park, K.-K.; Lee, S. S. Molecular mechanisms underlying chemopreventive activities of anti-inflammatory phytochemicals: down-regulation of COX-2 and iNOS through suppression of NF- κ B activation. *Mutat. Res.* **2001**, *480–481*, 243–268.
- (27) Rosen, G. M.; Tsai, P.; Pou, S. Mechanism of free-radical generation by nitric oxide synthase. *Chem. Rev.* **2002**, *102*, 1191–1199.
- (28) Park, Y. C.; Rimbach, G.; Saliou, C.; Valacchi, G.; Packer, L. Activity of monomeric, dimeric, and trimeric flavonoids on NO production, TNF- α secretion, and NF- κ B-dependent gene expression in RAW 264.7 macrophages. *FEBS Lett.* **2000**, *464*, 93–97.
- (29) Sheu, F.; Lai, H. H.; Yen, G. C. Suppression effect of soy isoflavones on nitric oxide production in RAW 264.7 macrophages. *J. Agric. Food Chem.* **2001**, *49*, 1767–1772.
- (30) Chiang, Y.-M.; Lo, C.-P.; Chen, Y.-P.; Wang, S.-Y.; Yang, N.-S.; Kuo, Y.-H.; Shyur, L.-F. Ethyl caffeate suppresses NF- κ B activation and its downstream inflammatory mediators, iNOS, COX-2, and PGE₂ in vitro or in mouse skin. *Br. J. Pharmacol.* **2005**, *146*, 352–363.
- (31) Kim, H. K.; Cheon, B. S.; Kim, Y. H.; Kim, S. Y.; Kim, H. P. Effects of naturally occurring flavonoids on nitric oxide production in the macrophage cell line RAW 264.7 and their structure–activity relationships. *Biochem. Pharmacol.* **1999**, *58*, 759–765.
- (32) Paul, A.; Pendreigh, R. H.; Plevin, R. Protein kinase C and tyrosine kinase pathway regulate lipopolysaccharide-induced nitric oxide synthase activity in RAW 264.7 murine macrophages. *Br. J. Pharmacol.* **1995**, *114*, 482–488.
- (33) Tero, J.; Karasawa, H.; Arai, H.; Nagao, A.; Susuki, T.; Takama, K. Peroxyl radical scavenging activity of caffeic acid and its related phenolic compounds in solution. *Biosci., Biotechnol., Biochem.* **1993**, *57*, 1204–1205.
- (34) Sakakibara, H.; Honda, Y.; Nakagawa, S.; Ashida, H.; Kanazawa, K. Simultaneous determination of all polyphenols in vegetables, fruit, and teas. *J. Agric. Food Chem.* **2003**, *51*, 571–581.
- (35) Owen, R. W.; Giacosa, A.; Hull, W. E.; Haubner, R.; Spiegelhalder, B.; Bartsch, H. The antioxidant/anticancer potential of phenolic compounds isolated from olive oil. *Eur. J. Cancer* **2000**, *36*, 1235–1247.
- (36) Yang, C. S.; Landau, J. M.; Huang, M. T.; Newmark, H. L. Inhibition of carcinogenesis by dietary polyphenolic compounds. *Annu. Rev. Nutr.* **2001**, *21*, 381–406.
- (37) Rice-Evans, C. A.; Miller, N. J. Antioxidant activities of flavonoids as bioactive components of food. *Biochem. Soc. Trans.* **1996**, *24*, 790–795.

Received for review October 2, 2007. Revised manuscript received December 26, 2007. Accepted December 27, 2007. We thank the National Science Council for financial support (NSC92-2313-B-002-022).

JF072922S